



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

### DIOPHANTINE ANALYSIS.

---

109. Proposed by HARRY S. VANDIVER, Bala, Pa.

If  $m+n+1$  is a prime integer, show that  $m! \times n! - (-1)^{\frac{1}{2}(3m-n)}$  is divisible by  $m+n+1$ . For instance,  $6! \times 4! - (-1)^7$  is divisible by 11.

110. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.

Prove the results stated in the foot-note at the end of §4 in the article by L. E. Dickson in the October number of the MONTHLY.

111. Proposed by HARRY S. VANDIVER, Bala, Pa.

Show that: If a rational, integral polynomial of the  $n$ th degree in  $x$ , becomes a prime for more than  $n$  values of  $x$ , then it cannot be resolved into rational factors. (Test with the expression  $x^4 + 2x^3 + x^2 + 2x + 1$ .)

### AVERAGE AND PROBABILITY.

---

135. Proposed by L. C. WALKER, A. M., Graduate Student, Leland Stanford Jr. University, Cal.

If the line joining two points taken at random in the surface of a given circle be the diagonal of a square, the chance that the square lies wholly within the circle is  $2 - 4/\pi$ .

---

## NOTES.

---

Prof. Lon C. Walker is doing postgraduate work in Leland Stanford Jr. University. F.

Prof. Harry S. Vandiver is taking a course of mathematics in the University of Pennsylvania. F.

On December 15, the University of Klausenberg will celebrate the hundredth anniversary of the birth of Johann Bolyai. F.

Dr. F. Schottky of Marburg has been named to succeed the late L. Fuchs as Professor in the University of Berlin. Dr. K. Hensel of Berlin has been named Professor in the University of Marburg. D.

Dr. Joseph Swain, President of the University of Indiana since 1893, and formerly Professor of Mathematics in Indiana and Stanford Universities, was installed as President of Swarthmore College on November 15. D.

In the reconstitution of the University of London, complete courses of study in the various faculties of the University College have been established.